






Revised Simplex method: Introduction Benefit of revised simplex method is clearly comprehended in case of large LP problems. In simplex method the entire simplex tableau is updated while a small part of it is used. The revised simplex method uses exactly the same steps as those in simplex method. The only difference occurs in the details of computing the entering variables and departing variable.
























Finding Dual of a LP problem	
Primal	Dual
Maximization	Minimization
Minimization	Maximization
ith variable	ith constraint
j th constraint	j th variable
$x_i > 0$	Inequality sign of i th Constraint: ≤ if dual is maximization ≥ if dual is minimization






Finding Dual of a LP problem...contd. Note: Before finding its dual, all the constraints should be transformed to 'less-than-equal-to' or 'equal-to' type for maximization problem and to 'greater-than-equal-to' or 'equal-to' type for minimization problem. It can be done by multiplying with -1 both sides of the constraints, so that inequality sign gets reversed.







Dual Simplex Method

Simplex Method verses Dual Simplex Method

- Simplex method starts with a nonoptimal but feasible solution where as dual simplex method starts with an optimal but infeasible solution.
- Simplex method maintains the feasibility during successive iterations where as dual simplex method maintains the optimality.

D Nagesh Kumar, IISc

LP_5: Revised Simplex, Dual etc

Dual Simplex Method: Iterative steps

Steps involved in the dual simplex method are:

- All the constraints (except those with equality (=) sign) are modified to 'less-than-equal-to' sign. Constraints with greater-than-equal-to' sign are multiplied by -1 through out so that inequality sign gets reversed. Finally, all these constraints are transformed to equality sign by introducing required slack variables.
- Modified problem, as in step one, is expressed in the form of a simplex tableau. If all the cost coefficients are positive (i.e., optimality condition is satisfied) and one or more basic variables have negative values (i.e., non-feasible solution), then dual simplex method is applicable.

D Nagesh Kumar, IISc

LP_5: Revised Simplex, Dual etc

Dual Simplex Method: Iterative steps...contd.

- Selection of exiting variable: The basic variable with the highest negative value is the exiting variable. If there are two candidates for exiting variable, any one is selected. The row of the selected exiting variable is marked as pivotal row.
- Selection of entering variable: Cost coefficients, corresponding to all the negative elements of the pivotal row, are identified. Their ratios are calculated i.e.,

$$ratio = \left(\frac{Cost \ Coefficients}{Elements \ of \ pivotal \ row}\right)$$

The column corresponding to minimum ratio is identified as the pivotal column and associated decision variable is the entering variable.

21

D Nagesh Kumar, IISc

LP_5: Revised Simplex, Dual etc

Dual Simplex Method: Iterative steps...contd.

- Pivotal operation: Pivotal operation is exactly same as in the case of simplex method, considering the pivotal element as the element at the intersection of pivotal row and pivotal column.
- Check for optimality: If all the basic variables have nonnegative values then the optimum solution is reached. Otherwise, Steps 3 to 5 are repeated until the optimum is reached.

2:

D Nagesh Kumar, IISc

LP_5: Revised Simplex, Dual etc

Dual Simplex Method: An Example

Consider the following problem:

Minimize
$$Z = 2x_1 + x_2$$

subject to $x_1 \ge 2$
 $3x_1 + 4x_2 \le 24$
 $4x_1 + 3x_2 \ge 12$
 $-x_1 + 2x_2 \ge 1$

23

D Nagesh Kumar, IISc

LP_5: Revised Simplex, Dual etc


Dual Simplex Method: An Example...contd.


After converting them to 'less than or equal to' type constraints and introducing the slack variables the problem is reformulated with equality constraints as follows:


24


D Nagesh Kumar, IISc


LP_5: Revised Simplex, Dual etc





























Comparative discussion between new methods and Simplex method

1. Both Khatchian's ellipsoid method and Karmarkar's projective scaling method have been shown to be polynomial time algorithms.

Time required for an LP problem of size n is at most anb, where a and b are two positive numbers.

2. Simplex algorithm is an exponential time algorithm in solving LP problems.

Time required for an LP problem of size n is at most c2n, where c is a positive number


Comparative discussion between new methods and Simplex method 3. For a large enough n (with positive a, b and c), $c2^n > an^b$. The polynomial time algorithms are computationally superior to exponential algorithms for large LP problems. 4. However, the rigorous computational effort of Karmarkar's projective scaling method, is not economical for 'not-so-large' problems. D Nagesh Kumar, IISc LP_5: Revised Simplex, Dual etc


Karmarkar's projective scaling method • Also known as Karmarkar's interior point LP algorithm • Starts with a trial solution and shoots it towards the optimum solution • LP problems should be expressed in a particular form


LP_5: Revised Simplex, Dual etc

LP_5: Revised Simplex, Dual etc

D Nagesh Kumar, IISc







LP_5: Revised Simplex, Dual etc

D Nagesh Kumar, IISc

LP formulation problem Two types of crops can be grown in a particular irrigation area each year. Each unit quantity of yield of crop. A produced can be sold for a price $P_{\rm A}$ and requires $W_{\rm A}$ units of water, $L_{\rm A}$ units of land, $F_{\rm A}$ units of fertilizer and $H_{\rm A}$ units of labour. Similarly, yield from crop B can be sold at a unit price of $P_{\rm B}$ and requires $W_{\rm B}$, $L_{\rm B}$, $F_{\rm B}$, and $H_{\rm B}$ units of water, land, fertilizer and labour respectively per unit of crop. Assume that the available quantity of water, land, fertilizer and labour are known and equal $W_{\rm L}$, $F_{\rm F}$ H respectively. Structure a LP model for estimating the quantities of each of the two crops that should be produced in order to maximize total income. Decision variables: X_A and X_B - Quantity of yield from crops A and B respectively Objective Function: $P_A X_A + P_B X_B$ Subject to: Water availability constraint $W_A X_A + W_B X_B \le W$ Land availability constraint $L_A \ X_A + \ L_B \ X_B \le L$ $F_A X_A + F_B X_B \leq F$ Fertilizer availability constraint $H_A X_A + H_B X_B \le H$ Labour availability constraint Non-negativity constraints $X_A \ge 0$ and $X_B \ge 0$

D Nagesh Kumar, IISc